

Analysis of Anomalous Interactions With Heavy Leptons at ep , e^+e^- and pp Colliders

A. T. Tasci* and A. T. Alan†

Abant Izzet Baysal University, Department of Physics, 14280 Bolu, Turkey

Abstract

We consider possible production of heavy leptons via anomalous interactions at future ep colliders (THERA and LHeC), e^+e^- colliders (ILC and CLIC) and pp collider CERN LHC. The production, backgrounds and signatures of heavy leptons are analyzed. We obtain the upper mass limits of 800 GeV at LHeC, 450 GeV at ILC and 650 GeV at LHC for optimal choices of relevant parameters.

PACS numbers: 12.60.-i, 13.66.De, 14.60.-z

*Electronic address: tasci.a@ibu.edu.tr

†Electronic address: alan.a@ibu.edu.tr

I. INTRODUCTION

Many models extending the standard theory of quarks and leptons predict the existence of new generations of fermions. New heavy leptons play an important role in the search for extensions of the Standard Model (SM) and any signal for the production of such fermions will play a milestone role in the discovery of new physics. Many analysis have been done for the production of these particles at future e^-e^+ [1, 2, 3, 4], at hadron [5, 6, 7, 8] and also at ep colliders [9, 10, 11, 12]. The experimental upper bounds for the heavy lepton masses were found to be 44 GeV by OPAL [13], 46 GeV by ALEPH [14] and 90 GeV by H1 [15] Collaborations. Hence, these leptons could be detected at future high energy colliders as their masses could be as high as 1 TeV. In this work, we analyze possible production and decay processes of new heavy leptons via some anomalous interactions in ep , e^-e^+ and pp collisions at future ep colliders THERA [16] and LHeC [17], at linear colliders ILC [18] and CLIC [19], at the CERN Large Hadron Collider (LHC) [20]. The main parameters of these colliders are given in Table I.

In the SM, Flavor Changing Neutral Current (FCNC) processes receive contributions from only higher order corrections. Here we offer the following effective Lagrangian having magnetic moment type operators to describe the interactions of the heavy leptons by ordinary ones;

$$\mathcal{L}_{\text{eff}} = \frac{iek_\gamma}{\Lambda} L \sigma_{\mu\nu} q^\nu l A^\mu + \frac{g}{2 \cos \theta_W} L \left[\gamma_\mu (c_v - c_a \gamma_5) + \frac{i \sigma_{\mu\nu} q^\nu}{\Lambda} \kappa_Z \right] l Z^\mu + h.c., \quad (1)$$

where κ_γ and κ_Z are the anomalous magnetic dipole moment factors, c_v and c_a are the corresponding anomalous non-diagonal Z couplings which are zero in the SM, q is the momentum of the exchanged gauge boson, θ_W is the Weinberg angle, e and g denote the gauge couplings relative to $U(1)$ and $SU(2)$ symmetries respectively, A^μ and Z^μ the fields of the photon and Z boson and Λ is the new physics scale. In the numerical calculations we have taken c_v and c_a at the order of SM values.

Heavy leptons decay through the neutral current processes $L \rightarrow \gamma l$ and $L \rightarrow Z l$ via anomalous couplings in Eq. (1), where l denotes one of the ordinary charged leptons (e, μ, τ). The branching ratios (BR) for these processes would be around 33% for each channel. Neglecting ordinary lepton masses the decay widths are obtained as,

$$\Gamma(L \rightarrow l \gamma) = \frac{\alpha \kappa_\gamma^2 m_L^3}{2 \Lambda^2}, \quad (2)$$

$$\Gamma(L \rightarrow lZ) = \frac{\alpha(m_L^2 - M_Z^2)^2}{16\Lambda^2 m_L^3 M_Z^2 \sin^2 \theta_W \cos^2 \theta_W} \left[\kappa_Z^2 M_Z^4 + 2M_Z^2 \left[\Lambda^2(c_v^2 + c_a^2) + (\kappa_Z^2 m_L^2 - 3c_v \kappa_Z \Lambda m_L) \right] + \Lambda^2(c_v^2 + c_a^2)m_L^2 \right], \quad (3)$$

where α is the electromagnetic coupling constant, M_Z and m_L refer to masses of Z boson and decaying lepton, respectively.

The only background processes related to the signal reaction $ep \rightarrow LqX$ are $eq \rightarrow e\gamma q$ and $eq \rightarrow eZq$. In Table II we have presented the decay widths $\Gamma(L \rightarrow \gamma e)$ and $\Gamma(L \rightarrow Ze)$ for a wide range of m_L . As seen from this table the branching ratio $\text{BR}(L \rightarrow \gamma e)$ is extremely small compared with $\text{BR}(L \rightarrow Ze)$, hence the background from γe channel can be safely ignored.

II. PRODUCTION OF HEAVY LEPTONS IN ep COLLISIONS

Using Eq. (1), the differential cross section for the subprocess $eq \rightarrow Lq$, through the t channel mediated by γ and Z is obtained as;

$$\begin{aligned} \frac{d\hat{\sigma}(eq \rightarrow Lq)}{d\hat{t}} &= \frac{2\kappa_\gamma^2 e_q^2 \pi \alpha^2}{\Lambda^2 \hat{s}^2 \hat{t}} \left\{ (2\hat{s} + \hat{t})m_L^2 - 2\hat{s}(\hat{s} + \hat{t}) - m_L^4 \right\} \\ &+ \frac{\pi \alpha^2}{8\Lambda^2 \hat{s}^2 \sin^4 \theta_W \cos^4 \theta_W [(\hat{t} - M_Z^2)^2 + M_Z^2 \Gamma_Z^2]} \left\{ 2\kappa_Z \Lambda c_v^l (c_a^{q2} + c_v^{q2})(m_L^2 - \hat{t})m_L \hat{t} \right. \\ &+ 4\Lambda c_v^q c_a^l (\Lambda c_v^l - m_L \kappa_Z)(m_L^2 - 2\hat{s} - \hat{t})\hat{t} \\ &- \Lambda^2(c_a^{q2} + c_v^{q2})(c_a^{l2} + c_v^{l2}) [(2\hat{s} + \hat{t})m_L^2 - 2\hat{s}^2 - 2\hat{s}\hat{t} - \hat{t}^2] \\ &\left. - \kappa_Z^2(c_a^{q2} + c_v^{q2}) [m_L^4 - (2\hat{s} + \hat{t})m_L^2 + 2\hat{s}(\hat{s} + \hat{t})] \hat{t} \right\} \\ &+ \frac{\kappa_\gamma e_q \pi \alpha^2 (\hat{t} - M_Z^2)}{\Lambda^2 \hat{s}^2 \sin^2 \theta_W \cos^2 \theta_W [(\hat{t} - M_Z^2)^2 + M_Z^2 \Gamma_Z^2]} \left\{ \kappa_Z c_v^q (m_L^4 - (2\hat{s} + \hat{t})m_L^2 + 2\hat{s}(\hat{s} + \hat{t})) \right. \\ &\left. + \Lambda m_L (c_a^l c_a^q (m_L^2 - 2\hat{s} - \hat{t}) + c_v^l c_v^q (\hat{t} - m_L^2)) \right\}, \quad (4) \end{aligned}$$

where e_q is quark charge in units of e , Γ_Z is the decay width of mediator Z . The total production cross section is obtained by the integration of differential cross section over the parton distributions in the proton as;

$$\sigma(ep \rightarrow LqX) = \int_{x_{min}}^1 dx f_q(x, Q^2) \int_{t_{min}}^{t_{max}} \frac{d\hat{\sigma}}{d\hat{t}} d\hat{t}, \quad (5)$$

where $x_{min} = m_L^2/s$, $\hat{t}_{min} = -(\hat{s} - m_L^2)$ and $\hat{t}_{max} = 0$. For the parton distribution functions $f_q(x, Q^2)$ we have used the CTEQ5 parametrization [21] providing the dependence

on momentum transfer which have been taken as $Q = m_L$ and for illustration, values of $\kappa_\gamma = \kappa_Z = 0.02$ have been taken for anomalous magnetic moment couplings.

We give the production cross sections for the signal as functions of the heavy lepton masses, m_L , in Fig. 1 for $\sqrt{s}=1$ and 1.4 TeV. We applied an optimal cut of $|M_{Ze} - m_L| < 50$ GeV for the considered heavy lepton mass range, in order to contain all signal events that are smeared by the experimental resolution. Fig. 2 shows the p_T distributions of the final state particles for the relevant background process $eq \rightarrow eZq$ in ep collisions. These p_T distributions have peaks around 50 GeV and suppressed at higher values. Fig. 3 shows the invariant mass distributions of Ze system, after a cut of $p_T^{e,j} > 50$ GeV.

In Tables III and IV we present the production and total background cross sections depending on the heavy lepton masses for suitable ranges at THERA and LHeC. The number of signal events per year can easily be estimated from the integrated luminosities of the colliders. The branching ratios BR_1 and BR_2 refer to $BR(L \rightarrow Ze)$ and $BR(Z \rightarrow e^+e^-, \mu^+\mu^-)$, respectively. As shown in Table IV production of heavy leptons provides a clean signature with small backgrounds. According to Poisson statistics, the statistical significance (SS) of the signal for the discovery at 95% confidence level obeys the relation,

$$\frac{S}{\sqrt{S+B}} \geq 3, \quad (6)$$

where S and B are the numbers of signal and background events, respectively. LHeC, with an integrated luminosity of 10^4 pb^{-1} yields 12 events per year for 800 GeV leptons, hence a good place to search for heavy leptons. On the other hand THERA seems capable for only about five 100 GeV leptons with $SS \simeq 2$ which is not sufficient for observation.

III. SINGLE AND PAIR PRODUCTION OF HEAVY LEPTONS IN e^+e^- COLLISIONS

A. Single Production

The single production of heavy leptons L , in e^-e^+ collisions occur through the s and t channels via the process $e^-e^+ \rightarrow L^-e^+$ represented by four Feynman diagrams, two for Z mediation plus two for photon mediation. The corresponding total cross section is obtained

as follows;

$$\begin{aligned}
\frac{d\sigma}{dt} = & \frac{\pi\alpha^2}{8\Lambda^2 s^2 \sin^4 \theta_W \cos^4 \theta_W [(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2][(t - M_Z^2)^2 + M_Z^2 \Gamma_Z^2]} \\
& \times \left\{ 2\Lambda \kappa_Z c_v s t (t - s) m_L [c_a^2 (7(s + t) - 6m_L^2) + c_v^2 (s + t - 2m_L^2)] - \kappa_Z^2 (c_a^2 + c_v^2) \right. \\
& \times s t [(s + t) m_L^4 - 2(s^2 + t^2) m_L^2 + 2(s^3 + t^3)] + \Lambda^2 [(c_a^4 + c_v^4) (2(s^4 - t^2 s^2 + t^4) \\
& - (s + t) (2s^2 - 3st + 2t^2) m_L^2) - 2c_a^2 c_v^2 ((s - 2t)(2s - t)(s + t) m_L^2 \\
& - 2(s^4 - 3t^2 s^2 + t^4))] + M_Z^2 [2\kappa_Z^2 (c_a^2 + c_v^2) (2m^4 - 2(s + t) m_L^2 + (s + t)^2) s t \\
& + 2\kappa_Z c_v \Lambda (s - t) m_L [c_a^2 (3s^2 + 3t^2 + 8st - 3(s + t) m_L^2) + c_v^2 (s^2 + t^2 - (s + t) m_L^2)] \\
& - 2\Lambda^2 (c_a - c_v)^2 (c_a + c_v)^2 (s^3 + t^3 - (s^2 + t^2) m_L^2) - (M_Z^2 + \Gamma_Z^2) \\
& \times [\kappa_Z^2 (c_a^2 + c_v^2) (s + t) (m_L^4 - (s + t) m_L^2 + 2st) + \Lambda^2 (c_a - c_v)^2 (c_a + c_v)^2 \\
& \times ((s + t) m_L^2 - s^2 - t^2)] \left. \right\} - \frac{2\pi\alpha^2 \kappa_\gamma^2}{\Lambda^2 s^3 t} \left\{ (s + t) m_L^4 - 2(s^2 + t^2) m_L^2 + 2(s^3 + t^3) \right\} \\
& + \frac{\pi\alpha^2 \kappa_\gamma M_Z \Gamma_Z}{\Lambda^2 s^2 \sin^2 \theta_W \cos^2 \theta_W [(t - M_Z^2)^2 + M_Z^2 \Gamma_Z^2]} \left\{ \kappa_Z c_v m_L^4 - 2\Lambda (c_a^2 + c_v^2) m_L^3 \right. \\
& - 2\kappa_Z c_v m_L^2 s + \Lambda (c_a^2 (3s + 2t) + c_v^2 (s + 2t)) m_L + \kappa_Z c_v (2s - t)(s + t) \left. \right\} \\
& + \frac{\pi\alpha^2 \kappa_\gamma M_Z \Gamma_Z}{\Lambda^2 s^2 \sin^2 \theta_W \cos^2 \theta_W [(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2]} \left\{ \kappa_Z c_v (m_L^4 - 2t m_L^2 - (s - 2t)(s + t)) \right. \\
& + \Lambda ((c_a^2 + c_v^2) (2m_L^2 - 2s - t) - 2c_a^2 t) m_L \left. \right\} \quad (7)
\end{aligned}$$

To compare the linear colliders, we display the total cross sections as functions of the heavy lepton masses in Fig. 4 by assuming $\Lambda = m_L$ and taking the anomalous coupling parameters as $\kappa_\gamma = \kappa_Z = 0.02$. Here, we applied the same cut, $|M_{Ze} - m_L| < 50$ GeV, as in ep collisions. Fig. 5 shows the p_T distributions of the final state particles at three different linear colliders. The character of this distribution is similar to that of ep collisions. In Fig. 6 we give the invariant mass distributions M_{Ze} , with cut $p_T^{e,j} > 50$ GeV.

In Tables V, VI and VII, we presented the signal and total background cross sections for the single production of heavy lepton depending on its mass m_L , for 0.5, 1 and 3 TeV energy e^-e^+ colliders, respectively. As seen from these tables, the SS values are high enough for observability limit, for mass values up to the center of mass energies of the colliders.

In the single production case we expect 777 events per year for 450 GeV leptons, 903 events for 900 GeV leptons and 810 events for 2750 GeV leptons at 0.5, 1 and 3 TeV, respectively.

B. Pair Production

Pair production of heavy leptons via anomalous couplings occur through the t -channel flavor changing neutral current process $e^-e^+ \rightarrow L^-L^+$. The differential cross section has the form;

$$\begin{aligned}
\frac{d\sigma_{tot.}}{dt} = & \frac{\pi\alpha^2\kappa_\gamma^2}{\Lambda^2s^2t^2} \left\{ 2m_L^8 - 4tm_L^6 + t(4s+3t)m_L^4 - 2t^2(4s+t)m_L^2 + t^2(2s+t)^2 \right\} \\
& + \frac{\pi\alpha^2}{16M_Z^4\Lambda^4s^2\sin^4\theta_W\cos^4\theta_W[(t-M_Z^2)^2+M_Z^2\Gamma_Z^2]} \\
& \times \left\{ \Lambda^4(c_a^2+c_v^2)m_L^4[(m_L^2-t)^2+4M_Z^2s] \right. \\
& - 2\Lambda^2M_Z^2\kappa_Z^2c_v^2m_L^2t[m_L^4+t(t+2s-2m_L^2)] \\
& + M_Z^4[2\Lambda^4[(c_a^4+c_v^4)(m_L^4-2(2s+t)m_L^2+2s^2+t^2+2st) \\
& + 2c_a^2c_v^2(3m_L^4-2(2s+3t)m_L^2+2s^2+3t^2+6st)] \\
& - 2\kappa_Z^2\Lambda^2[4(c_a^2+c_v^2)s(m_L^4+st+t^2)+(c_a^2+2c_v^2)(m_L^6-2tm_L^4+t^2m_L^2) \\
& - 10c_v^2t m_L^2] + \kappa_Z^2(2m_L^8-4tm_L^6+tm_L^4(4s+3t) \\
& \left. - 2t^2(4s+t)m_L^2+t^2(2s+t)^2)] \right\} \\
& + \frac{\pi\alpha^2\kappa_\gamma^2(t-M_Z^2)}{2M_Z^2\Lambda^4s^2t\sin^2\theta_W\cos^2\theta_W[(t-M_Z^2)^2+M_Z^2\Gamma_Z^2]} \\
& \times \left\{ \kappa_Z^2M_Z^2[2m_L^8-4tm_L^6+t(4s+3t)m_L^4-2t^2(4s+t)m_L^2 \right. \\
& + t^2(2s+t)^2] - 2\Lambda^2M_Z^2m_L^2[(c_a^2+c_v^2)(m_L^4+t^2-2tm_L^2) \\
& \left. + st(2c_a^2-c_v^2)] - c_v^2\Lambda^2tm_L^2[m_L^4-2tm_L^2+(2s+t)t] \right\} \quad (8)
\end{aligned}$$

The total cross sections as functions of heavy lepton masses m_L are displayed in Fig. 7 for three options.

Signal and total background cross sections depending again on the heavy lepton masses, are presented in Tables VIII, IX and X at 0.5, 1 and 3 TeV, respectively.

We applied an initial cut on the electron and jet transverse momentum $p_T^{e,j} > 50$ GeV for the background analysis. Fig. 8 show the p_T distributions of the final state particles at the colliders. The distribution of invariant mass M_{Ze} is presented in Fig. 9 at $\sqrt{s}=0.5, 1$ and 3 TeV.

From Table X it is seen that the number of signal events could be as high as 8×10^4 for 1250 GeV leptons at $\sqrt{s} = 3$ TeV.

IV. PRODUCTION OF HEAVY LEPTONS AT THE CERN LHC

In pp collisions heavy lepton production occurs via parton level subprocess $q\bar{q} \rightarrow Ll$, which are γ and Z exchanged s channel FCNC reactions.

The form of the differential cross section is as follows;

$$\begin{aligned}
\frac{d\hat{\sigma}(q\bar{q} \rightarrow Ll)}{d\hat{t}} = & \frac{2e_q^2\kappa_\gamma^2\pi\alpha^2}{\Lambda^2\hat{s}^3} \left\{ (\hat{s} + 2\hat{t})m_L^2 - m_L^4 - 2\hat{t}(\hat{s} + \hat{t}) \right\} \\
& + \frac{\pi\alpha^2}{8\Lambda^2\hat{s}^2 \sin^4\theta_W \cos^4\theta_W [(\hat{s} - M_Z^2)^2 + M_Z^2\Gamma_Z^2]} \\
& \times \left\{ \kappa_Z^2(c_a^{q2} + c_v^{q2})\hat{s}((\hat{s} + 2\hat{t})m_L^2 - 2\hat{t}(\hat{s} + \hat{t}) - m_L^4) \right. \\
& + 2\kappa_Z\Lambda\hat{s}m_L(c_v(c_a^{q2} + c_v^{q2})(\hat{s} - m_L^2) - 2c_ac_a^q c_v^q(m_L^2 - \hat{s} - 2\hat{t})) \\
& + \Lambda^2[(c_a^2 + c_v^2)(c_a^{q2} + c_v^{q2})(\hat{s}^2 + 2\hat{t}^2 + 2\hat{s}\hat{t} - (\hat{s} + 2\hat{t})m_L^2) \\
& \left. + 4c_ac_v c_a^q c_v^q \hat{s}(\hat{s} + 2\hat{t} - m_L^2)] \right\} \\
& + \frac{e_q\pi\alpha^2\kappa_\gamma M_Z\Gamma_Z}{\Lambda^2\hat{s}^2 \sin^2\theta_W \cos^2\theta_W [(\hat{s} - M_Z^2)^2 + M_Z^2\Gamma_Z^2]} \\
& \times \left\{ \kappa_Z c_v^q(m_L^4 - (\hat{s} + 2\hat{t})m_L^2 + 2\hat{t}(\hat{s} + \hat{t})) \right. \\
& \left. + \Lambda m_L(c_ac_a^q(m_L^2 - \hat{s} - 2\hat{t}) + c_v c_v^q(m_L^2 - \hat{s})) \right\} \quad (9)
\end{aligned}$$

The total production cross section is obtained by the integration over the parton distributions as;

$$\sigma(q\bar{q} \rightarrow Ll) = \int_{m_L^2/s}^1 d\tau \int_\tau^1 \frac{dx}{x} [f_{q/p}(x, Q^2)f_{\bar{q}}(x/\tau, Q^2) + f_q(x/\tau, Q^2)f_{\bar{q}}(x, Q^2)]\hat{\sigma}(\hat{s}) \quad (10)$$

where $\hat{s} = \tau s$. As in the ep collisions, here we use the CTEQ5 parametrization with $Q = m_L$ and we take $\kappa_\gamma = \kappa_Z = 0.02$ and $\Lambda = m_L$.

We give the production cross sections for the signal as functions of the heavy lepton masses, m_L , in Fig. 10 for the center of mass energy of $\sqrt{s} = 14$ TeV. We applied an optimal cut of $|M_{Ze} - m_L| < 50$ GeV for the considered heavy lepton mass range. Fig. 11 shows the p_T distribution of the final state particles for the relevant background process $q\bar{q} \rightarrow eZl$ in pp collisions. This p_T distribution has a peak around 50 GeV and suppressed at higher values. Fig. 12 shows the invariant mass distributions of Ze system, after a cut of $p_T^{e,j} > 50$ GeV.

In Table XI, we present the production cross sections and the number of signal events depending on the heavy lepton masses for a suitable range. The branching ratios BR_1 and BR_2 refer to $BR(L \rightarrow Ze)$ and $BR(Z \rightarrow e^+e^-, \mu^+\mu^-)$, respectively. Production of heavy leptons provides a clean signature, as shown in this table. Taking into account the SS, LHC can observe heavy leptons with masses up to about 650 GeV. For 650 GeV leptons we expect 11 events per year at the LHC.

We have used the high energy package CompHEP for calculations of background cross sections reported in this study [22].

V. CONCLUSION

In conclusion, the present work gives an analysis of possible production of heavy leptons via anomalous interactions in ep , e^+e^- and pp collisions. It is shown that, after kinematical cuts a statistical significance of $SS \geq 3$ can be achieved and for anomalous magnetic moment factors of $\kappa_\gamma = \kappa_Z = 0.02$, heavy leptons with masses about 800 GeV at LHeC, 450 GeV at ILC and 650 GeV at the LHC can be observed. Among the others ILC seems the most convenient place to search for heavy leptons as providing a high event number of 8×10^4 . LHeC and LHC are also good places for searching heavy leptons besides the other major high energy experiments. Hence, these collider options seem to be capable of probing new physics in the case of anomalous interactions are valid as being an underlying theory.

Acknowledgments

This study was partially supported by Abant Izzet Baysal University Research fund (2005.03.02.216).

-
- [1] F. M. L. Almeida, J. H. Lopes, J. A. Martins Simoes and C. M. Porto, Phys. Rev. D **44**, 2836 (1991).
 - [2] F. M. L. Almeida, J. H. Lopes, J. A. Martins Simoes, P. P. Queiroz Filho and A. J. Ramalho, Phys. Rev. D **51**, 5990 (1995).

- [3] F. M. L. Almeida, Y. A. Coutinho, J. A. Martins Simoes and M. A. B. do Vale, Phys. Rev. D **63**, 075005 (2001).
- [4] F. M. L. Almeida, Y. A. Coutinho, J. A. Martins Simoes, S. Wolck and M. A. B. do Vale, Eur. Phys. J. C **30**, 327 (2003).
- [5] P. H. Frampton, D. Ng, M. Sher and Y. Yuan, Phys. Rev. D **48**, 3128 (1993).
- [6] Y. A. Coutinho, J. A. Martins Simoes, C. M. Porto and P. P. Queiroz Filho, Phys. Rev. D **57**, 6975 (1998).
- [7] J. E. Cieza Montalvo and P. P. de Queiroz Filho, Phys. Rev. D **66**, 055003 (2002).
- [8] J. E. Cieza Montalvo and M. D. Tonasse, Nucl. Phys. B **623**, 325 (2002).
- [9] F. M. L. Almeida, J. A. Martins Simoes and A. J. Ramalho, Nucl. Phys. B **347** (1990) 537.
- [10] T. G. Rizzo, Phys. Lett. B **188**, 95 (1987).
- [11] A. T. Alan, A. T. Tasci and O. Cakir, Acta Phys. Polon. B **35**, 2199 (2004).
- [12] A. T. Alan, A. T. Tasci and N. Karagoz, Mod. Phys. Lett. A **21**, 1869 (2006).
- [13] M. Z. Akrawy *et al.* [OPAL Collaboration], Phys. Lett. B **240**, 250 (1990).
- [14] D. Decamp *et al.* [ALEPH Collaboration], Phys. Rept. **216**, 253 (1992).
- [15] T. Ahmed *et al.* [H1 Collaboration], Phys. Lett. B **340**, 205 (1994).
- [16] H. Abramowicz *et al.* [TESLA-N Study Group], DESY-01-011.
- [17] J. B. Dainton, M. Klein, P. Newman, E. Perez and F. Willeke, DESY-06-006, arXiv:hep-ex/0603016.
- [18] International Linear Collider-Technical Review Report, ILC-TRC/2003 Report (2003).
- [19] E. Accomando *et al.* [CLIC Physics Working Group], “Physics at the CLIC multi-TeV linear collider”.
- [20] D. Boussard *et al.*, [LHC Study Group], CERN-AC-95-05-LHC (1995).
- [21] H. L. Lai *et al.* [CTEQ Collaboration], Eur. Phys. J. C **12**, 375 (2000).
- [22] A. Pukhov *et al.*, arXiv:hep-ph/9908288.

TABLE I: Main parameters of ep and e^+e^- and pp colliders, L^{int} denotes the integrated luminosity for one year.

ep Colliders	E_e (TeV)	E_p (TeV)	$\sqrt{s_{ep}}$ (TeV)	$L_{ep}^{int}(\text{pb}^{-1})$
THERA	0.25	1	1	40
LHeC	0.07	7	1.4	10^4
e^+e^- Colliders	E_{e^+} (TeV)	E_{e^-} (TeV)	$\sqrt{s_{e^+e^-}}$ (TeV)	$L_{e^+e^-}^{int}(\text{pb}^{-1})$
ILC	0.25	0.25	0.5	10^5
CLIC	0.5	0.5	1.0	10^5
CLIC	1.5	1.5	3.0	10^5
pp Collider	E_p (TeV)	E_p (TeV)	$\sqrt{s_{pp}}$ (TeV)	$L_{pp}^{int}(\text{pb}^{-1})$
LHC	7	7	14	10^4

TABLE II: Decay widths of heavy leptons.

m_L (GeV)	$\Gamma(L \rightarrow \gamma e)(\text{GeV})$	$\Gamma(L \rightarrow Ze)(\text{GeV})$	$\Gamma_{\text{Total}}(\text{GeV})$
250	7.30×10^{-4}	3	3
500	1.46×10^{-3}	21	21
750	2.19×10^{-3}	72	72
1000	2.92×10^{-3}	171	171
1250	3.65×10^{-3}	333	333
1500	4.38×10^{-3}	576	576
1750	5.11×10^{-3}	914	914
2000	5.84×10^{-3}	1365	1365
2250	6.57×10^{-3}	1943	1943
2500	7.30×10^{-3}	2666	2666

TABLE III: The signal and background cross sections and SS depending on the heavy lepton masses for THERA with $\sqrt{s} = 1$ TeV.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	$\sigma_B(\text{pb})$	$S/\sqrt{S+B}$
100	12.26	4.05	0.13	4.53×10^{-5}	2
150	8.21	2.71	0.09	2.54×10^{-4}	2
200	5.50	1.82	0.06	2.39×10^{-4}	2

TABLE IV: The signal and background cross sections and SS depending on the heavy lepton masses for LHeC with $\sqrt{s} = 1.4$ TeV.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	$\sigma_B(\text{pb})$	$S/\sqrt{S+B}$
200	9.43	3.11	0.194	4.75×10^{-4}	32
400	2.80	0.93	0.031	2.28×10^{-4}	17
600	0.68	0.22	0.007	8.69×10^{-5}	9
800	0.11	0.04	0.001	2.19×10^{-5}	3

TABLE V: The signal and background cross sections and SS depending on the heavy lepton masses at $\sqrt{s} = 0.5$ TeV for single production.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	$\sigma_B(\text{pb})$	$S/\sqrt{S+B}$
100	4.63	1.53	0.0504	1.28×10^{-2}	63
200	3.90	1.29	0.0424	5.10×10^{-3}	62
300	2.88	0.95	0.0314	2.05×10^{-3}	54
400	1.51	0.50	0.0165	4.83×10^{-4}	40

TABLE VI: The signal and background cross sections and SS depending on the heavy lepton masses at $\sqrt{s} = 1$ TeV for single production.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	σ_B (pb)	$S/\sqrt{S+B}$
100	4.89	1.61	0.053	1.43×10^{-2}	65
300	4.30	1.42	0.047	4.60×10^{-3}	65
500	3.51	1.16	0.038	2.36×10^{-3}	60
700	2.35	0.78	0.026	1.04×10^{-3}	50
900	0.83	0.27	0.009	1.81×10^{-4}	30

TABLE VII: The signal and background cross sections and SS depending on the heavy lepton masses at $\sqrt{s} = 3$ TeV for single production.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	σ_B (pb)	$S/\sqrt{S+B}$
250	4.74	1.56	0.052	6.37×10^{-3}	68
750	4.44	1.46	0.048	3.55×10^{-3}	67
1250	3.91	1.29	0.043	3.04×10^{-3}	63
1750	3.12	1.03	0.034	2.44×10^{-3}	56
2250	2.06	0.68	0.022	1.43×10^{-3}	46
2750	0.74	0.25	0.008	3.06×10^{-4}	28

TABLE VIII: The signal and background cross sections and SS depending on the heavy lepton masses at $\sqrt{s} = 0.5$ TeV for pair production.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	σ_B (pb)	$S/\sqrt{S+B}$
100	3.95	1.30	0.0430	3.22×10^{-2}	50
150	3.76	1.24	0.0409	9.20×10^{-3}	58
200	3.83	1.26	0.0417	1.78×10^{-3}	63
240	2.79	0.92	0.0304	1.98×10^{-2}	43

TABLE IX: The signal and background cross sections and SS depending on the heavy lepton masses at $\sqrt{s} = 1$ TeV for pair production.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	σ_B (pb)	$S/\sqrt{S+B}$
100	4.51	1.49	0.0491	9.92×10^{-3}	64
200	4.48	1.48	0.0488	4.91×10^{-3}	67
300	5.72	1.89	0.0623	7.31×10^{-3}	75
400	9.44	3.12	0.1028	4.24×10^{-3}	99

TABLE X: The signal and background cross sections and SS depending on the heavy lepton masses at $\sqrt{s} = 3$ TeV for pair production.

m_L (GeV)	σ (pb)	$\sigma \times \text{BR}_1$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2$ (pb)	σ_B (pb)	$S/\sqrt{S+B}$
250	4.72	1.56	0.0514	2.11×10^{-3}	70
500	6.12	2.02	0.0666	3.32×10^{-4}	81
750	15.03	4.96	0.1637	4.11×10^{-4}	128
1000	39.49	13.03	0.4300	1.93×10^{-4}	207
1250	77.64	25.62	0.8455	1.32×10^{-5}	291

TABLE XI: The signal and background cross sections and SS depending on the heavy lepton masses for the LHC ($\sqrt{s} = 14$ TeV).

m_L (GeV)	$\sigma \times 10^{-2}$ (pb)	$\sigma \times \text{BR}_1 \times 10^{-2}$ (pb)	$\sigma \times \text{BR}_1 \times \text{BR}_2 \times 10^{-4}$ (pb)	$\sigma_B \times 10^{-4}$ (pb)	$S/\sqrt{S+B}$
150	9.22	3.04	10.0	4.96	8
250	6.80	2.24	7.4	4.05	7
350	4.58	1.51	5.0	2.80	6
450	2.86	0.95	3.1	1.84	4
550	1.70	0.56	1.9	1.14	3
650	0.98	0.32	1.1	0.65	3




FIG. 1: The total cross sections as function of the heavy lepton masses for lepton-hadron colliders THERA ($\sqrt{s} = 1$ TeV) and LHeC ($\sqrt{s} = 1.4$ TeV).




FIG. 2: p_T distributions of the background for lepton-hadron colliders THERA and LHeC.




FIG. 3: The invariant mass distributions of the Ze system for the background for lepton-hadron colliders THERA and LHeC.




FIG. 4: The total cross sections as function of the heavy lepton masses for linear colliders ILC ($\sqrt{s}=0.5$ TeV) and CLIC ($\sqrt{s}=1$ and 3 TeV) for single production.




FIG. 5: p_T distribution of the background for linear colliders ILC and CLIC for single production.




FIG. 6: The invariant mass distribution of the Ze system for the background for linear colliders ILC and CLIC for single production.




FIG. 7: The total cross sections as function of the heavy lepton masses for linear colliders ILC and CLIC for pair production.




FIG. 8: p_T distribution of the background for linear colliders ILC and CLIC for pair production.




FIG. 9: The invariant mass distribution of the Ze system for the background for linear colliders ILC and CLIC for pair production.




FIG. 10: The total cross sections as function of the heavy lepton masses for LHC ($\sqrt{s}=14$ TeV).




FIG. 11: p_T distribution of the background for LHC.




FIG. 12: The invariant mass distribution of the Ze system for the background for LHC.